Emergencies in Migrant and Refugee Health 2015-2018 (letter)

E. Kalavsky (Erich Kalavsky)², G. Herdics (Georgy Herdics)¹, J. Vallova (Jana Vallova)¹, S. Subramaniam (Selvaraj Subramaniam)², J. Bydzovsky (Jan Bydzovsky)¹, J. Ridosko (Jaroslav Ridosko)³, T. Simonek (Tomas Simonek)², M. Palun (Miroslav Palun)², P. Vansac (Peter Vansac)¹, A. Naddour (Annemarie Naddour)², J. Suvada (Jose Suvada)²

¹ SEUC Refugee and Migrant Health Program, Bratislava, SK		Original Article
 ² MSc/Nutrition Program Kuala Lumpur, Malaysia ³ University Hospital Trencin, SK 	, SK	
E-mail address:		
houseoffamily@hotmail.com		
Reprint address:		
Erich Kalavsky		
House of Family		
PnomPenh, KH		
Source: Clinical Social Work and Health Intervention Pages: 56 – 57	Volume: 9 Cited references	Issue: 3
1 uges. 50 – 57	Clieu rejerences.	7
Reviewers:		
Daria Kimuli – pechacova.daria@gmail.com		
Mageswari Rajoo – m-rajoo@hotmail.com		
Key words:		
Refugees. Migrants. Emergencies.		
Publisher:		
International Society of Applied Preventive Medicine i-ga	p	
CSWHI 2018; 9(3): 56 – 57; DOI 10.22359/cswhi_9_3_06 © 2018 Clinical	Social Work and Health	1 Intervention

Sir,

Several papers in Clinical Social Work and Health Interventions have been related to spectrum of either communicable or non-communicable diseases (1-4). However none of them report emergency situations. They have been suspected to be related to PTSD, asthma, coronary heart disease, hypertension and diabetes; all diseases which will exacerbate with acute stress; conditions refugees and migrants must tolerate during their stressful journey to EU. Table 1 shows the commonest emergencies in various Health Checkpoints and Health Centers near Refugee Camps, for example in Hegyeshalom (HU), Dobova (SLO), Veria (GR), Lesbos (GR). The commonest emergency was diabetic coma, followed by status asthmaticus, acute bleeding during birth and hypertension crisis. Surprisingly, no death was recorded in any of those events despite of the large number of migrants coming via the Balkan Route.

Table 1: RList of commonest emergency events
for acute care

Event	No.	Deaths
Birth bleeding	12	0
Bleeding wounds	8	0
Tentamen Suicidi	2	0
Diabetic coma	16	0
Myocardial infarction	7	0
Status asthmaticus	6	0

References

- MOLNAROVA K, ONDOVA P, ZOLLER K, ADAMCOVA A et al. (2017) Low Number of Neuroinfections in Migrants to Greece from Syria and Iraq Comparison to Migrants from Burundi and Congo to Rwanda. Clinical Social Work and Health Intervention Vol. 8 No. 2 2017.
- BELOVICOVA M, LISKOVA A, HRIN-DOVA T et al. (2017) Low Prevalence of MRSA in Physiotherapy and Gym Facilities in a Greek Refugee Camp. Clinical Social Work and Health Intervention Vol. 8 No. 2 2017.
- KHALED I, ALI HAJJ P, KRCMERY V, KALATOVA D, GALLOVA A, BAK T, SCHUMMAN I, DUDOVA Z, ZOLLER K, REITH B, OBTULOVIC M, OLAH M, HRINDOVA T, BELOVICOVA M (2017) Spectrum of infections in physiotherapy and rehabilitation ward for war victims and veterans from Iraq and Syria (short note). Clinical Social Work and Health Intervention. 8. 26-28. 10.22359/cswhi 8 2 05.
- HAJJ PA et al. (2016) Are migrants from Middle East carriers of multi-resistant bacteria? Clinical Social Work and Health Intervention Vol. 7 No. 3 2016 2.